

IN THE CLAIMS:

This listing of claims will replace all prior versions and listings of claims in the application:

Claims 1-7 (cancel)

8. (New) A pharmaceutical composition obtained from the green or mature fruits of *Roystonea regia* comprising:

a mixture of primary fatty acids with 8 to 28 carbon atoms, the fatty acid selected from the group consisting of caprylic acid (C8:0), capric acid (C10:0), lauric acid (C12:0), miristic acid (C14:0), palmitic acid (C16:0), palmitoleic acid (C16:1), stearic acid (C18:0), oleic acid (C18:1), linoleic acid, and linolenic acid; and

a mixture of esters of the fatty acids;

wherein free fatty acids are enriched from esters hydrolysis.

9. (New) The pharmaceutical composition according to claim 1 comprising:

Caprylic acid (C8:0)	< 3.0 %
Capric acid (C10:0)	< 3.0 %
Lauric acid (C12:0)	3.0 - 40.0 %
Miristic acid (C14:0)	4.0 - 15.0 %
Palmitic acid (C16:0)	10.0 - 80.0 %
Palmitoleic acid (C16:1)	1.5 - 20.0 %
Stearic acid (C18:0)	0.1 - 5.0 %
Oleic acid (C18:1)	3.0 - 50.0 %

U.S. Application No.: New
PRELIMINARY AMENDMENT

Attorney Docket: 4082.003

10. (New) A method for the obtention of the pharmaceutical composition obtained from *Roystonea regia* according to claim 8 comprising:

drying, grounding, and sieving the *Roystonea regia* fruits, and

separating of the extract from other components through a solid/liquid extraction in organic solvents,

wherein the organic solvents are chosen from hydrocarbons of 5 to 8 carbon atoms, alcohols of 1 to 3 carbon atoms, and mixture thereof, with or without a previous basic hydrolysis using hydroxides or alkalis.

11. (New) The method according to the claim 10 wherein the drying of *Roystonea regia* fruits is at a temperature between 10 and 100 °C for a time ranging from 1 to 1000 hours, and

the grounding is to obtain a particle size < 6000 µm;

the extraction is for 1 to 50 h at a temperature from 0 to 70 °C.

12. (New) The method according to claim 10 wherein the hydrolysis includes the use of alkaline- hydroxides or alkaline-earthen hydroxides and organic for the basic hydrolysis, specifically those of low molecular weight selected from the group consisting of sodium, potassium, calcium or ammonium hydroxides.

U.S. Application No.: New
PRELIMINARY AMENDMENT

Attorney Docket: 4082.003

13. (New) The method according to claim 10 wherein the hydrocarbons are selected from the group consisting of pentane, hexane, heptane, or octane.

14. (New) The method according to claim 10 wherein the alcohol is selected from the group consisting of methanol, ethanol, n-propanol, and 2-propanol.

15. (New) A method for treating and/or to prevent BPH, prostatitis, alopecia and hirsutism comprising administering a medication comprising the pharmaceutical composition according to claim 8.

16. (New) The method according to claim 15 wherein the pharmaceutical composition is with or without a saponification.

17. (New) The method of claim 15 wherein the medicament is administered at daily doses from 50 to 1000 mg.

18. (New) The method of claim 15 wherein the medicament is administered at doses between 150 and 1000 mg.

19. (New) The method of claim 15 wherein the medicament is administered as solid oral forms (capsules, soft-gel capsules, tablets), liquids (emulsions), suppositories, tinctures, lotions, or shampoos of local action.